

Many Nodal Domains in Random Regular Graphs

Theo McKenzie

with Shirshendu Ganguly, Sidhanth Mohanty, and Nikhil Srivastava
University of California, Berkeley

Online Asymptotic Geometric Analysis Seminar

04/05/2022

Quantum Chaos vs Integrable Systems

A central theme in mathematical physics is the dichotomy between integrable systems and quantum chaotic systems.

Quantum Chaos vs Integrable Systems

A central theme in mathematical physics is the dichotomy between integrable systems and quantum chaotic systems.

Integrable systems: We expect spectral fluctuations to be Poisson, and eigenfunctions to be localized in phase space.

Quantum chaotic systems: We expect spectral fluctuations to be those of large random matrices, and eigenfunctions to be equidistributed in phase space.

Behavior of Eigenfunctions

- [Shnirelman 1974] Quantum Ergodicity Theorem: A Riemannian manifold with ergodic geodesic flow is such that almost all high energy eigenfunctions of the Laplacian are equidistributed.

Behavior of Eigenfunctions



- [Shnirelman 1974] Quantum Ergodicity Theorem: A Riemannian manifold with ergodic geodesic flow is such that almost all high energy eigenfunctions of the Laplacian are equidistributed.
- [Berry 1977] Random wave conjecture: in the high energy limit, statistics of eigenfunctions of surfaces with quantum dynamics behave like Gaussian random variables with covariance

$$E(\psi_n(x)\psi_n(y)) = J_0(k_n|x - y|).$$

- J_0 is the 0th order Bessel function and k_n^2 is the eigenvalue of ψ_n .

Discrete Graphs as a Model for Quantum Chaos

- [Kottos-Smilansky 1997,1999] Initiate using large regular graphs as a model for quantum chaos by examining the eigenvectors of the discrete Laplacian.
- Rather than take the high energy limit, we send the number of vertices to ∞ .



Discrete Graphs as a Model for Quantum Chaos

- [Kottos-Smilansky 1997,1999] Initiate using large regular graphs as a model for quantum chaos by examining the eigenvectors of the discrete Laplacian.
- Rather than take the high energy limit, we send the number of vertices to ∞ .
- [Anantharaman-Le Masson 2015] Prove a statement analogous to Shnirelman's Theorem for large regular graphs.

Discrete Graphs as a Model for Quantum Chaos

- [Kottos-Smilansky 1997,1999] Initiate using large regular graphs as a model for quantum chaos by examining the eigenvectors of the discrete Laplacian.
- Rather than take the high energy limit, we send the number of vertices to ∞ .
- [Anantharaman-Le Masson 2015] Prove a statement analogous to Shnirelman's Theorem for large regular graphs.
- [Elon 2008]: Formulated Berry's Conjecture for discrete graphs.

Nodal Domains

- How do we test Berry's conjecture?

Nodal Domains

- How do we test Berry's conjecture?
- One way is to choose a specific statistic, and then measure that statistic versus what is implied by Berry's conjecture.

Nodal Domains


- How do we test Berry's conjecture?
- One way is to choose a specific statistic, and then measure that statistic versus what is implied by Berry's conjecture.
- If the statistic matches Berry's conjecture, then it can be interpreted as a suggestion of the conjecture.

Nodal Domains

- How do we test Berry's conjecture?
- One way is to choose a specific statistic, and then measure that statistic versus what is implied by Berry's conjecture.
- If the statistic matches Berry's conjecture, then it can be interpreted as a suggestion of the conjecture.
- **Nodal domains** are a way to measure randomness in both continuous and discrete space.

Courant's Nodal Domain Theorem

- [Courant] The zero set of the k th smallest Dirichlet eigenfunction of the Laplacian on a smooth bounded domain in \mathbb{R}^d partitions it into at most k components.
- These components, known as nodal domains, have garnered significant attention in spectral geometry and mathematical physics.



A heat map of the 6th Dirichlet eigenfunction of the square.

Discrete Version


- There is also a well-studied theory of nodal domains on graphs.

Discrete Version

- There is also a well-studied theory of nodal domains on graphs.
- The **nodal domains** of a vector f on the vertices of a graph G are the maximal connected components of all the same sign.


Discrete Version

- The **nodal domains** of a vector f on the vertices of a graph G are the maximal connected components of all the same sign.




Discrete Version

- The **nodal domains** of a vector f on the vertices of a graph G are the maximal connected components of all the same sign.




Discrete Version

- The **nodal domains** of a vector f on the vertices of a graph G are the maximal connected components of all the same sign.




Discrete Version

- The **nodal domains** of a vector f on the vertices of a graph G are the maximal connected components of all the same sign.




Discrete Version

- The **nodal domains** of a vector f on the vertices of a graph G are the maximal connected components of all the same sign.




Berry's Conjecture to Many Nodal Domains

- **Claim:** In both continuous and discrete space, Berry's conjecture implies the existence of many nodal domains.




Berry's Conjecture to Many Nodal Domains

- **Claim:** In both continuous [Bogomolny and Schmit] and discrete space [Elon], Berry's conjecture implies the existence of many nodal domains.
- **Goal:** Show in the discrete model families of graphs with many nodal domains.



Adjacency Matrix

- Encode graphs through an adjacency matrix A , with rows/columns corresponding to the vertices, and placing a 1 where there is an edge.



$$\begin{bmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

- Note that as the matrix is symmetric, the eigenvalues are real.
- We can also consider the combinatorial Laplacian $D - A$, where D is the diagonal matrix of degrees.

Graph Nodal Domains

- [Fiedler 1975] The eigenvector of the k th smallest eigenvalue of the combinatorial Laplacian of a tree has exactly k nodal domains.

Graph Nodal Domains

- [Fiedler 1975] The eigenvector of the k th smallest eigenvalue of the combinatorial Laplacian of a tree has exactly k nodal domains.
- [Davies et al. 2000] For any graph, under general assumptions the k th eigenvector has at most k nodal domains.

Graph Nodal Domains

- [Fiedler 1975] The eigenvector of the k th smallest eigenvalue of the combinatorial Laplacian of a tree has exactly k nodal domains.
- [Davies et al. 2000] For any graph, under general assumptions the k th eigenvector has at most k nodal domains.
- [Berkolaiko 2008] For a graph on n vertices and $n + \ell - 1$ edges, the k th eigenvector of a Schrödinger operator of arbitrary potential has between $k - \ell$ and k nodal domains (generalizes the above two results).

Graph Nodal Domains


- [Fiedler 1975] The eigenvector of the k th smallest eigenvalue of the combinatorial Laplacian of a tree has exactly k nodal domains.
- [Davies et al. 2000] For any graph, under general assumptions the k th eigenvector has at most k nodal domains.
- [Berkolaiko 2008] For a graph on n vertices and $n + \ell - 1$ edges, the k th eigenvector of a Schrödinger operator of arbitrary potential has between $k - \ell$ and k nodal domains (generalizes the above two results).
- Question: Can we lower bound the number of nodal domains for graphs that have many edges?

Graph Nodal Domains

- [Fiedler 1975] The eigenvector of the k th smallest eigenvalue of the combinatorial Laplacian of a tree has exactly k nodal domains.
- [Davies et al. 2000] For any graph, under general assumptions the k th eigenvector has at most k nodal domains.
- [Berkolaiko 2008] For a graph on n vertices and $n + \ell - 1$ edges, the k th eigenvector of a Schrödinger operator of arbitrary potential has between $k - \ell$ and k nodal domains (generalizes the above two results).
- Question: Can we lower bound the number of nodal domains for graphs that have many edges?
- What about random graphs?

Erdős-Rényi Graphs

- For a graph on n vertices, include each of the $\binom{n}{2}$ potential edges independently with probability p . This distribution is denoted by $G(n, p)$.



Sampled from $G(10, 0.5)$

Results for Erdős-Rényi Graphs

- [Dekel-Lee-Linial 2011] For a $G(n, p)$ graph for fixed $p \in (0, 1)$, with high probability every eigenvector of the **adjacency matrix** besides the first has all but $O(1)$ vertices in two nodal domains.

Results for Erdős-Rényi Graphs

- [Dekel-Lee-Linial 2011] For a $G(n, p)$ graph for fixed $p \in (0, 1)$, with high probability every eigenvector of the **adjacency matrix** besides the first has all but $O(1)$ vertices in two nodal domains.
- [Arora-Bhaskara 2011] For a $G(n, p)$ graph with $p \in (n^{-1/20}, 1)$ with high probability every eigenvector besides the first has exactly two nodal domains.

Results for Erdős-Rényi Graphs

- [Dekel-Lee-Linial 2011] For a $G(n, p)$ graph for fixed $p \in (0, 1)$, with high probability every eigenvector of the **adjacency matrix** besides the first has all but $O(1)$ vertices in two nodal domains.
- [Arora-Bhaskara 2011] For a $G(n, p)$ graph with $p \in (n^{-1/20}, 1)$ with high probability every eigenvector besides the first has exactly two nodal domains.
- [Rudelson 2017] With high probability, every vertex neighbors the nodal domain of the opposite sign.

Results for Erdős-Rényi Graphs

- [Dekel-Lee-Linial 2011] For a $G(n, p)$ graph for fixed $p \in (0, 1)$, with high probability every eigenvector of the **adjacency matrix** besides the first has all but $O(1)$ vertices in two nodal domains.
- [Arora-Bhaskara 2011] For a $G(n, p)$ graph with $p \in (n^{-1/20}, 1)$ with high probability every eigenvector besides the first has exactly two nodal domains.
- [Rudelson 2017] With high probability, every vertex neighbors the nodal domain of the opposite sign.
- [H. Huang-Rudelson 2020] For bulk eigenvectors with $p \in (n^{-c}, 1)$ and edge eigenvectors for fixed $p \in (0, 1)$ the nodal domains are approximately the same size.

Random Regular Graphs


- Erdős-Rényi graphs are too dense to have more than two nodal domains for $p \geq n^{-c}$.

Random Regular Graphs

- Erdős-Rényi graphs are too dense to have more than two nodal domains for $p \geq n^{-c}$.
- Dekel, Lee, and Linial also observed that according to simulations, the k th eigenvector of a randomly selected d -regular graph on n vertices has a number of nodal domains that increases with k .

Random Regular Graphs

- Erdős-Rényi graphs are too dense to have more than two nodal domains for $p \geq n^{-c}$.
- Dekel, Lee, and Linial also observed that according to simulations, the k th eigenvector of a randomly selected d -regular graph on n vertices has a number of nodal domains that increases with k .




Random Regular Graphs

- Erdős-Rényi graphs are too dense to have more than two nodal domains for $p \geq n^{-c}$.
- Dekel, Lee, and Linial also observed that according to simulations, the k th eigenvector of a randomly selected d -regular graph on n vertices has a number of nodal domains that increases with k .
- Our question: can we prove nontrivial structure of nodal domains for a randomly selected d -regular graph, for fixed d .

Random Regular Graphs

- Choose a graph from the set of graphs on n vertices that are d -regular. We denote this distribution by $G(n, d)$.



- Sampled from $G(20,3)$

Quantum Chaos through Nodal Domains

- Showing many nodal domains for random d -regular graphs would give evidence towards Berry's conjecture on discrete graphs.

Result

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

Result


[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- The limiting spectral measure of random regular graphs means there are at least $cd^{-3/2}n$ eigenvalues in this interval for large enough n and some fixed $c > 0$.

Result

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.



- The limiting spectral measure of random regular graphs means there are at least $cd^{-3/2}n$ eigenvalues in this interval for large enough n and some fixed $c > 0$.



Result

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- The limiting spectral measure of random regular graphs means there are at least $cd^{-3/2}n$ eigenvalues in this interval for large enough n and some fixed $c > 0$.



Result

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- The limiting spectral measure of random regular graphs means there are at least $cd^{-3/2}n$ eigenvalues in this interval for large enough n and some fixed $c > 0$.
- This is the first result lower bounding the number of nodal domains for random regular graphs.

Result

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- The limiting spectral measure of random regular graphs means there are at least $cd^{-3/2}n$ eigenvalues in this interval for large enough n and some fixed $c > 0$.
- This is the first result lower bounding the number of nodal domains for random regular graphs.
- For regular graphs, A and the combinatorial Laplacian $L = D - A$ have the same eigenvectors, so we can rephrase the result as concerning the high energy eigenvectors of the Laplacian.

Result


[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- What we prove is that the number of **singleton** nodal domains (nodal domains of one vertex) is $\Omega(n/\text{polylog}(n))$.

Result

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o_n(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- What we prove is that the number of **singleton** nodal domains (nodal domains of one vertex) is $\Omega(n/\text{polylog}(n))$.



Outline

- We split into cases based on whether the eigenvector is **localized** or **delocalized** (whether the mass of the eigenvector is well spread or not).

- **Definition:** an eigenvector ψ is **delocalized** if for fixed $\epsilon, \delta > 0$,
 $|\{v \in V | \psi^2(v) \geq \epsilon/n\}| \geq \delta n$.

Outline

- We split into cases based on whether the eigenvector is **localized** or **delocalized** (whether the mass of the eigenvector is well spread or not).
- **Definition:** an eigenvector ψ is **delocalized** if for fixed $\epsilon, \delta > 0$,
$$|\{v \in V | \psi^2(v) \geq \epsilon/n\}| \geq \delta n.$$
- If the eigenvector is delocalized, we can use the proximity of an eigenvector of a random regular graph to a Gaussian distribution.

Outline

- We split into cases based on whether the eigenvector is **localized** or **delocalized** (whether the mass of the eigenvector is well spread or not).
- **Definition:** an eigenvector ψ is **delocalized** if for fixed $\epsilon, \delta > 0$,
$$|\{v \in V \mid \psi^2(v) \geq \epsilon/n\}| \geq \delta n.$$
- If the eigenvector is delocalized, we can use the proximity of an eigenvector of a random regular graph to a Gaussian distribution.
- If the eigenvector is localized, then we can argue using the local structure of random regular graphs.

Outline

- We split into cases based on whether the eigenvector is **localized** or **delocalized** (whether the mass of the eigenvector is well spread or not).

- **Definition:** an eigenvector ψ is **delocalized** if for fixed $\epsilon, \delta > 0$,
$$|\{v \in V \mid \psi^2(v) \geq \epsilon/n\}| \geq \delta n.$$

- If the eigenvector is delocalized, we can use the proximity of an eigenvector of a random regular graph to a Gaussian distribution.

- If the eigenvector is localized, then we can argue using the local structure of random regular graphs.

The Gaussian Wave

- Recall the covariance in Berry's conjecture is

$$E(\psi_n(x)\psi_n(y)) = J_0(k_n|x - y|).$$

The Gaussian Wave

- Recall the covariance in Berry's conjecture is

$$E(\psi_n(x)\psi_n(y)) = J_0(k_n|x - y|).$$


- An analogous statement on graphs would mean that entries are Gaussian and $E(\psi(u)\psi(v))$ should depend on the distance between u and v .

The Gaussian Wave

- Recall the covariance in Berry's conjecture is

$$E(\psi_n(x)\psi_n(y)) = J_0(k_n|x - y|).$$

- An analogous statement on graphs would mean that entries are Gaussian and $E(\psi(u)\psi(v))$ should depend on the distance between u and v .
- Almost everywhere, a random regular graph will locally look like a tree (they converge to the infinite tree T_d in the sense of Benjamini and Schramm).




The Gaussian Wave

- Recall the covariance in Berry's conjecture is

$$E(\psi_n(x)\psi_n(y)) = J_0(k_n|x - y|).$$

- An analogous statement on graphs would mean that entries are Gaussian and $E(\psi(u)\psi(v))$ should depend on the distance between u and v .
- Almost everywhere, a random regular graph will locally look like a tree (they converge to the infinite tree T_d in the sense of Benjamini and Schramm).



- Therefore, first we can try to understand covariance in the infinite d -regular tree.


The Gaussian Wave

- [Elon 2009] There is a unique joint Gaussian distribution on the vertices of T_d that
 1. Has unit variance in each entry
 2. Is automorphism invariant
 3. Satisfies the eigenvector equation at each vertex.

- This joint Gaussian is called the **Gaussian wave**.


Gaussian Wave

1. Has unit variance in each entry
2. Is automorphism invariant
3. Satisfies the eigenvector equation at each vertex



Gaussian Wave


1. Has unit variance in each entry
2. Is automorphism invariant
3. Satisfies the eigenvector equation at each vertex



$$\lambda X_0 = \sum_{v \sim 0} X_v$$

Gaussian Wave


1. Has unit variance in each entry
2. Is automorphism invariant
3. Satisfies the eigenvector equation at each vertex



$$\lambda X_0 = \sum_{v \sim 0} X_v$$
$$\lambda X_0^2 = \sum_{v \sim 0} X_v X_0$$

Gaussian Wave


1. Has unit variance in each entry
2. Is automorphism invariant
3. Satisfies the eigenvector equation at each vertex



$$\lambda X_0 = \sum_{v \sim 0} X_v$$
$$\lambda X_0^2 = \sum_{v \sim 0} X_v X_0$$
$$\lambda E(X_0^2) = \sum_{v \sim 0} E(X_v X_0)$$

Gaussian Wave

1. Has unit variance in each entry
2. Is automorphism invariant
3. Satisfies the eigenvector equation at each vertex



$$\lambda X_0 = \sum_{v \sim 0} X_v$$


$$\lambda X_0^2 = \sum_{v \sim 0} X_v X_0$$

$$\lambda E(X_0^2) = \sum_{v \sim 0} E(X_v X_0)$$

$$\lambda = dE(X_1 X_0)$$

Gaussian Wave

1. Has unit variance in each entry
2. Is automorphism invariant
3. Satisfies the eigenvector equation at each vertex



$$\lambda X_0 = \sum_{v \sim 0} X_v$$

$$\lambda X_0^2 = \sum_{v \sim 0} X_v X_0$$

$$\lambda E(X_0^2) = \sum_{v \sim 0} E(X_v X_0)$$

$$\lambda = d E(X_1 X_0)$$

$$E(X_0 X_1) = \lambda/d$$

Convergence to the Gaussian wave

- This means that if Berry's conjecture were true in the discrete setting, the covariances should be those of the Gaussian wave.



Convergence to the Gaussian wave

- This means that if Berry's conjecture were true in the discrete setting, the covariances should be those of the Gaussian wave.
- [Backhausz-Szegedy 2019] Prove a form of Berry's conjecture.

Backhausz and Szegedy

- For any $R, \epsilon > 0$, for large enough n , with probability at least $1 - \epsilon$, a random regular graph has the following property:
- For any eigenvector of the adjacency matrix, sample a vertex uniformly at random.
- The distribution of values in the R neighborhood of the vertex is at most ϵ in the weak topology from the distribution of the R -neighborhood in a multiple of the Gaussian wave.


Illustration of BS 19



The distribution of $\sqrt{n}(v_0, v_1, v_2, v_3)$ when the central vertex is selected uniformly at random is close to the distribution of the Gaussian wave $\sigma \cdot (X_0, X_1, X_2, X_3)$ for some $\sigma \in [0,1]$.


Localization

- For negative λ , there is a constant probability that entries in the Gaussian wave correspond to a singleton nodal domain.




Localization

- For negative λ , there is a constant probability that entries in the Gaussian wave correspond to a singleton nodal domain.




- However, the Gaussian wave our distribution is close to could have low variance (even variance 0). In this case, the eigenvector distribution's proximity to the Gaussian wave does not imply that there are many nodal domains.
- In this case, the eigenvector is **localized**.

Localization




If the eigenvector entries are close to a Gaussian with nonzero variance, then the proximity to the Gaussian wave implies many nodal domains

Localization




If the eigenvector entries are close to a Gaussian with nonzero variance, then the proximity to the Gaussian wave implies many nodal domains




If the entries are close to a Gaussian with variance close to zero, then there is no such implication, but we also know that most eigenvector entries are concentrated around 0.


Localization




Localization




Localization



Localization



Localization



Outline

- We split into cases based on whether the eigenvector is **localized** or **delocalized** (whether the mass of the eigenvector is well spread or not).

- **Definition:** an eigenvector ψ is **delocalized** if for fixed $\epsilon, \delta > 0$,
 $|\{v \in V \mid \psi^2(v) \geq \epsilon/n\}| \geq \delta n$.

- If the eigenvector is delocalized, we can use the proximity of an eigenvector of a random regular graph to a Gaussian distribution. ✓

- If the eigenvector is localized, then we can argue using the local structure of random regular graphs.

Outline

- We split into cases based on whether the eigenvector is **localized** or **delocalized** (whether the mass of the eigenvector is well spread or not).

- **Definition:** an eigenvector ψ is **delocalized** if for fixed $\epsilon, \delta > 0$,
 $|\{v \in V | \psi^2(v) \geq \epsilon/n\}| \geq \delta n$.

- If the eigenvector is delocalized, we can use the proximity of an eigenvector of a random regular graph to a Gaussian distribution. ✓

- If the eigenvector is localized, then we can argue using the local structure of random regular graphs.

Green's Function Bound

- It remains to solve the problem when the eigenvector is localized.
- We use another connection between the infinite tree and the random regular graph.

- [Bauerschmidt-Huang-Yau 2019, Huang-Yau 2021⁺] With high probability, the adjacency matrix A_G of a random regular graph G sampled from $G(n, d)$ is such that for any $z \in \mathbb{C}$ with $\Im(z) \geq \log^c n/n$


$$(z - A_G)^{-1}_{u,v} \approx (z - A_{T_d})^{-1}_{u^*,v^*}$$

where $u^*, v^* \in T_d$ and the graph distance of u^* and v^* is the graph distance of u and v .

Function Bound Continued


- We want to use the Green's function to analyze the structure of one eigenvector. However, because we must have $\Im(z) \geq \log^c n/n$, we cannot separate one eigenvector from its closest $\log^c n$ neighbors.

z 's Distance from the Real Line




- As the average distance between eigenvalues is $1/n$, for $(z - A)^{-1}$ to “focus” on one eigenvector, we should have $\Im(z) = \Theta(1/n)$.

z 's Distance from the Real Line



- As the average distance between eigenvalues is $1/n$, for $(z - A)^{-1}$ to "focus" on one eigenvector, we should have $\Im(z) = \Theta(1/n)$.



- If $\Im(z) \geq \log^c n/n$, then we cannot separate the desired eigenspace from the closest $\log^c n$ eigenspaces.

Infinity norm

- Nevertheless, the Green's function bound implies a nearly optimal bound on the ∞ -norm.
- [Bauerschmidt-Huang-Yau '19, Huang-Yau '21⁺] Corollary: With high probability, any eigenvector ψ of the adjacency matrix of a $G(n, d)$ graph satisfies

$$\|\psi\|_{\infty} \leq \log^{c/2} n / \sqrt{n}.$$

- Using this, then further analyzing properties of the localized vector, we can prove our result.

Proof for very negative eigenvalues

- Here we present a simpler version of our ideas, that only uses eigenvector delocalization, and no other properties of random regular graphs.

Proof for very negative eigenvalues continued

- Assume that ψ is an eigenvector of eigenvalue $\lambda \leq -d + 1 - \epsilon$.

Proof for very negative eigenvalues continued

- Assume that ψ is an eigenvector of eigenvalue $\lambda \leq -d + 1 - \epsilon$.
- Consider a vertex u that is **not** a singleton nodal domain such that $|\psi(u)| \geq \frac{1}{2\sqrt{n}}$.

Proof for very negative eigenvalues continued

- Assume that ψ is an eigenvector of eigenvalue $\lambda \leq -d + 1 - \epsilon$.
- Consider a vertex u that is **not** a singleton nodal domain such that $|\psi(u)| \geq \frac{1}{2\sqrt{n}}$.

$$\lambda\psi(u) = \sum_{v \sim u} \psi(v)$$

Proof for very negative eigenvalues continued

- Assume that ψ is an eigenvector of eigenvalue $\lambda \leq -d + 1 - \epsilon$.
- Consider a vertex u that is **not** a singleton nodal domain such that $|\psi(u)| \geq \frac{1}{2\sqrt{n}}$.

$$\lambda\psi(u) = \sum_{v \sim u} \psi(v)$$

- As u has at most $d - 1$ neighbors such that $\psi(u)\psi(v) \leq 0$, it must be the case that there is some neighbor v_1 of u such that

$$|\psi(v_1)| \geq \left(1 + \frac{\epsilon}{d-1}\right) |\psi(u)|$$

Proof for very negative eigenvalues continued

- Assume that ψ is an eigenvector of eigenvalue $\lambda \leq -d + 1 - \epsilon$.
- Consider a vertex u that is **not** a singleton nodal domain such that $|\psi(u)| \geq \frac{1}{2\sqrt{n}}$.

$$\lambda\psi(u) = \sum_{v \sim u} \psi(v)$$

- As u has at most $d - 1$ neighbors such that $\psi(u)\psi(v) \leq 0$, it must be the case that there is some neighbor v_1 of u such that

$$|\psi(v_1)| \geq \left(1 + \frac{\epsilon}{d-1}\right) |\psi(u)|$$

- If v_1 is not a singleton nodal domain, we can repeat this process.

Doing this k times we have v_k with $|\psi(v_k)| \geq \left(1 + \frac{\epsilon}{d-1}\right)^k |\psi(v_k)|$.

However, by the ∞ -norm bound, we must have $k \leq C \log \log n$, meaning we must reach a singleton nodal domain within this number of steps.

Proof for very negative eigenvalues continued

- Assume that ψ is an eigenvector of eigenvalue $\lambda \leq -d + 1 - \epsilon$.
- Consider a vertex u that is **not** a singleton nodal domain such that $|\psi(u)| \geq \frac{1}{2\sqrt{n}}$.

$$\lambda\psi(u) = \sum_{v \sim u} \psi(v)$$

- As u has at most $d - 1$ neighbors such that $\psi(u)\psi(v) \leq 0$, it must be the case that there is some neighbor v_1 of u such that

$$|\psi(v_1)| \geq \left(1 + \frac{\epsilon}{d-1}\right) |\psi(u)|$$

- If v_1 is not a singleton nodal domain, we can repeat this process.

Doing this k times we have v_k with $|\psi(v_k)| \geq \left(1 + \frac{\epsilon}{d-1}\right)^k |\psi(v_k)|$.

However, by the ∞ -norm bound, we must have $k \leq C \log \log n$, meaning we must reach a singleton nodal domain within this number of steps.

- By delocalization, there are many such starting vertices u , meaning there are many singleton nodal domains.

Small sets on random graphs

- With high probability, a $G(n, d)$ graph does not have eigenvalues below $-2\sqrt{d-1} - o_n(1)$, whereas this method works for $\lambda \leq -d + 1 - \epsilon$. Therefore, this is not useful for random regular graphs when $d \geq 5$.

Small sets on random graphs

- With high probability, a $G(n, d)$ graph does not have eigenvalues below $-2\sqrt{d-1} - o_n(1)$, whereas this method works for $\lambda \leq -d + 1 - \epsilon$. Therefore, this is not useful for random regular graphs when $d \geq 5$.
- To achieve our better bound, we work globally rather than locally.



Global Argument

- Consider the set of vertices u such that $|\psi(u)| \geq \epsilon/\sqrt{n}$ for some small ϵ . Namely, there is nontrivial eigenvector mass on the vertex.

Global Argument



- Consider the set of vertices u such that $|\psi(u)| \geq \epsilon/\sqrt{n}$ for some small ϵ . Namely, there is nontrivial eigenvector mass on the vertex.
- Assume that none of these vertices are singleton nodal domains. **Then we can remove an edge neighboring each of these vertices while only decreasing the Rayleigh quotient.**

Global Argument




- $\psi^T A \psi = \sum_{u \sim v} \psi(u) \psi(v) = \lambda.$

Global Argument



- $\psi^T A \psi = \sum_{u \sim v} \psi(u) \psi(v) = \lambda.$
- If u is a singleton nodal domain, then all terms corresponding to u in $\psi^T A \psi$ are negative.

Global Argument



- $\psi^T A \psi = \sum_{u \sim v} \psi(u) \psi(v) = \lambda.$
- If u is a singleton nodal domain, then all terms corresponding to u in $\psi^T A \psi$ are negative.
- However, if u is not a singleton nodal domain, then there is at least one term that is positive.

Subgraphs

- We first localize onto the set of vertices that contain the majority of the ℓ_2 mass in the eigenvector.

Subgraphs

- We first localize onto the set of vertices that contain the majority of the ℓ_2 mass in the eigenvector.
- If there are few vertices that are singleton nodal domains, we further localize onto vertices that are not singleton nodal domains.

Subgraphs

- We first localize onto the set of vertices that contain the majority of the ℓ_2 mass in the eigenvector.
- If there are few vertices that are singleton nodal domains, we further localize onto vertices that are not singleton nodal domains.
- We then delete all edges (u, v) such that $\psi(u)\psi(v) \geq 0$.


Subgraphs

- We first localize onto the set of vertices that contain the majority of the ℓ_2 mass in the eigenvector.
- If there are few vertices that are singleton nodal domains, we further localize onto vertices that are not singleton nodal domains.
- We then delete all edges (u, v) such that $\psi(u)\psi(v) \geq 0$.
- As there are no singleton nodal domains, we have deleted at least one edge from each remaining vertex.


Subgraphs

- We first localize onto the set of vertices that contain the majority of the ℓ_2 mass in the eigenvector.
- If there are few vertices that are singleton nodal domains, we further localize onto vertices that are not singleton nodal domains.
- We then delete all edges (u, v) such that $\psi(u)\psi(v) \geq 0$.
- As there are no singleton nodal domains, we have deleted at least one edge from each remaining vertex.
- We are left with a subgraph H of maximum degree $d - 1$ such that the vector projected onto this subgraph has Rayleigh quotient
$$\psi_H^T A_H \psi_H \leq \psi^T A \psi + \epsilon$$
for some small ϵ .


Process




Process




Process




Process



Process



Process



At no step did we significantly increase the Rayleigh quotient of ψ .
Therefore, $\psi_S^T A_S \psi_S \leq \lambda + \epsilon$.

This subgraph is **small** and **has maximum degree $d - 1$** .

Local Structure

- Once again, random regular graphs locally look like trees.

Local Structure

- Once again, random regular graphs locally look like trees.
- We are localized on a small set, so our graph is close to tree-like.

Local Structure

- Once again, random regular graphs locally look like trees.
- We are localized on a small set, so our graph is close to tree-like.
- Theorem [Kesten 1959]: The spectral radius of the adjacency matrix of a tree of degree at most Δ is at most $2\sqrt{\Delta - 1}$.


Local Structure

- Once again, random regular graphs locally look like trees.
- We are localized on a small set, so our graph is close to tree-like.
- Theorem [Kesten 1959]: The spectral radius of the adjacency matrix of a tree of degree at most Δ is at most $2\sqrt{\Delta - 1}$.
- **Claim:** Because the maximum degree is $d - 1$ and the graph is close to tree-like, the spectral radius cannot be more than $2\sqrt{d - 2} + \epsilon$.

Local Structure

- Once again, random regular graphs locally look like trees.
- We are localized on a small set, so our graph is close to tree-like.
- Theorem [Kesten 1959]: The spectral radius of the adjacency matrix of a tree of degree at most Δ is at most $2\sqrt{\Delta - 1}$.
- **Claim:** Because the maximum degree is $d - 1$ and the graph is close to tree-like, the spectral radius cannot be more than $2\sqrt{d - 2} + \epsilon$.
- This can be proven through proving that the quadratic form on this tree-like graph is a convex combination of the quadratic forms of $(1 + \delta)\psi A_T \psi$, where A_T is the adjacency matrix of a subtree.


Local Structure Continued



- The resulting graph is close to treelike and has maximum degree $d - 1$. Therefore, the spectral radius is $\leq 2\sqrt{d - 2} + \epsilon$

$$\lambda = \psi^T A \psi \approx \psi_H^T A_H \psi_H \geq -2\sqrt{d - 2} - \epsilon$$

Local Structure Continued



- The resulting graph is close to treelike and has maximum degree $d - 1$. Therefore, the spectral radius is $\leq 2\sqrt{d - 2} + \epsilon$
$$\lambda = \psi^T A \psi \approx \psi_H^T A_H \psi_H \geq -2\sqrt{d - 2} - \epsilon$$
- By the ∞ -norm bound, we must have deleted **many** edges.

Recap

- As is the case with previous nodal domain results, we require bounds on the delocalization of our eigenvector, even though we prove a lower bound instead of an upper bound on the number of nodal domains.

Recap

- As is the case with previous nodal domain results, we require bounds on the delocalization of our eigenvector, even though we prove a lower bound instead of an upper bound on the number of nodal domains.
- We even do this in the localized case.

In the Future

- We want to improve the parameters of our theorem.

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

In the Future

- We want to improve the parameters of our theorem.

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- To quantify the $1 - o(1)$ term, we would need to bound the rate of convergence of Backhausz-Szegedy to the Gaussian wave. This would involve analyzing the entropy of the limiting process.

In the Future

- We want to improve the parameters of our theorem.

[Ganguly-M-Mohanty-Srivastava] Fix $d \geq 3$ and $\alpha > 0$. Then with probability $1 - o(1)$, every eigenvector of the adjacency matrix of a $G(n, d)$ sampled graph with eigenvalue $\lambda \leq -2\sqrt{d-2} - \alpha$ has $\Omega(n/\text{polylog}(n))$ nodal domains.

- To quantify the $1 - o(1)$ term, we would need to bound the rate of convergence of Backhausz-Szegedy to the Gaussian wave. This would involve analyzing the entropy of the limiting process.
- To improve the bound on λ , perhaps we could utilize the lack of nodal domains of different sizes besides just singletons, or find other “holes” in the spectrum.

In the Future

- Another problem is if we were able to prove “no gaps” delocalization then we would show that the local structure is governed by the Gaussian wave.

In the Future

- Another problem is if we were able to prove “no gaps” delocalization then we would show that the local structure is governed by the Gaussian wave.
- This would predict that there are many nodal domains for all eigenvectors with eigenvalue $\leq 2\sqrt{d-1} - \epsilon$, and would also predict their distribution.

In the Future

- Another problem is if we were able to prove “no gaps” delocalization then we would show that the local structure is governed by the Gaussian wave.
- This would predict that there are many nodal domains for all eigenvectors with eigenvalue $\leq 2\sqrt{d-1} - \epsilon$, and would also predict their distribution.
- We expect there to be many nodal domains of small size, and then two large nodal domains that contain almost all vertices.

In the Future

- Another problem is if we were able to prove “no gaps” delocalization then we would show that the local structure is governed by the Gaussian wave.
- This would predict that there are many nodal domains for all eigenvectors with eigenvalue $\leq 2\sqrt{d-1} - \epsilon$, and would also predict their distribution.
- We expect there to be many nodal domains of small size, and then two large nodal domains that contain almost all vertices.

Claim: At least $\left(1 - \frac{5}{\sqrt{d}}\right)$ vertices lie in the largest negative and largest positive nodal domain.

In the Future

- Finally, it is worth mentioning that in the original Erdős-Rényi case, little is known about $G(n, p)$ for $p \leq n^{-1/20}$. Eldan, H. Huang and Rudelson suggested considering the critical level for connectivity $p = c \log n/n$ to see if for p at this level there are many nodal domains.

In the Future

- Finally, it is worth mentioning that in the original Erdős-Rényi case, little is known about $G(n, p)$ for $p \leq n^{-1/20}$. Eldan, H. Huang and Rudelson suggested considering the critical level for connectivity $p = c \log n/n$ to see if for p at this level there are many nodal domains.
- Linial suggested to consider the geometry of nodal domains. This includes classifying vertices by their distance to the boundary of the nodal domain.

Thank you!